30 pts

Due Monday, November 3rd, at the beginning of class. I will have zero patience for stapling during class time.

Use separate paper to do the work on this take-home test. Start a fresh sheet of paper to show work on #4. Use paper without lines. Use only one side of each sheet of paper. *I will not grade work written on the backs of pages*. Write clearly and make sure your pencil work is *dark*. It's a struggle for me to read faint print.

Let $f(x) = x^5 - x^4 - 5x^3 + 9x^2 - 16x - 60$. We'll say everything we can about this polynomial that's worth saying.

- 1. (3 pts) Describe the end behavior of the graph of f with a simple graphic.
- 2. (3 pts) Use Descartes' Rule of Signs to determine the *possible* number of positive and negative zeros of f.
- 3. (3 pts) Use the Rational Zeroes Theorem to determine the possible rational zeroes of f.
- 4. (3 pts) Informed by your work, above, *and maybe a graphing utility of some sort*, use synthetic division to find the zeros. Each time you find a zero, it *should* reduce (depress) the question by one degree. Each time you find a zero, you should thereafter be working with a *depressed polynomial* that is of lesser degree.
- 5. (3 pts) From you work, above, factor f over the real numbers. This will involve an irreducible quadratic factor.
- 6. (3 pts) From your work above, factor f over the complex numbers. This should split f into linear factors.
- 7. (3 pts) Give a rough sketch of f that shows all intercepts.
- 8. (3 pts) Sketch the graph of $R(x) = \frac{x^2 5x 6}{x^2 5x + 6}$. Show all asymptotes, intercepts and any holes.
- 9. (3 pts) The graph of $g(x) = \frac{x^3 9x^2 + 14x + 24}{x^3 9x^2 + 26x 24}$ differs from the graph of f, in #8, in only one small detail. Sketch the graph of g, showing all asymptotes, intercepts and holes.
- 10. (3 pts) Sketch the graph of $R(x) = \frac{x^3 4x^2 7x + 10}{x^2 x 6}$, showing all asymptotes, intercepts and holes.