

Test 1 #5e

$$y = \cot(\sec(x^2-5)) \implies$$


$$y' = -\csc^2(\sec(x^2-5)) (\sec(x^2-5) + \tan(x^2-5)) (2x)$$

$$\frac{d}{dx} [f(g(h(x)))] = \frac{df}{dg} \cdot \frac{dg}{dh} \cdot \frac{dh}{dx}$$

$$\textcircled{3} \lim_{x \rightarrow 5} (2x-7) = 3$$

Proof Let  $\epsilon > 0$  be given. Define  $\delta = \frac{\epsilon}{2}$   
Then any time  $0 < |x-5| < \delta$ , we have

$$|(2x-7) - 3| = |2x-10| = 2|x-5| < 2\delta = \epsilon \quad \square$$



Build the function

$$-4 \leq x < 1$$

$$m = \frac{-4-3}{-4-1} = \frac{-7}{-5} = \frac{7}{5}$$

$$y = \frac{7}{5}(x-1) + 3$$

$$x = 1, y = 1$$

$$1 < x < \infty$$

$$m = \frac{-2-5}{4-1} = \frac{-7}{3}$$

$$y = -\frac{7}{3}(x-1) + 5$$




FIGURE 1




FIGURE 2

Abs min  $f(a)$ , abs max  $f(d)$ ,  
loc min  $f(c)$ ,  $f(e)$ , loc max  $f(b)$ ,  $f(d)$





FIGURE 3

### §3.1 Optimization



① Absolute max on  $D$   
at  $x=c$  means  $f(c) \geq f(x)$   
 $\forall x \in D$

(for every  $x$  in the Domain)

Abs. Min:  $f(c) \leq f(x) \forall x \in D$ .

② Local max at  $c$  means  
 $f(c) \geq f(x) \forall x$  on an  
open interval containing  $c$ .  
Likewise local min,  
 $f(c) \leq f(x) \forall x \in (a,b)$

③ Extreme Value Theorem  
Every continuous function on  
a closed interval has an abs.  
min and an abs. max on  
the interval.

Fermat's Theorem - If  $f(c)$  is a max/min and  $f'(c) \exists$ , then  $f'(c) = 0$ .  
(Sometimes you can have a max where  $f'(c) \nexists$ )

---

Critical #s: Any  $x$  where  $f'(x) = 0$  or  $f'(x) \nexists$

---

Optimization Recipe on closed interval  $[a, b]$

- ① Critical #s,  $c$ . Find the  $f(c)$ 's
- ② Find  $f(a)$  &  $f(b)$
- ③ Biggest is max.

#45-56

Find Abs. Max & Min

(46)  $f(x) = -2x^3 + 54x + 5$  on  $[0, 4]$

$f'(x) = -6x^2 + 54 \stackrel{\text{SET}}{=} 0$

$6x^2 - 54 = 0$

$x^2 - 9 = 0$

$x = \pm 3$

$f(3) = -2(3)^3 + 54(3) + 5$

$= -54 + 162 + 5$

$= \boxed{221 = f(3)}$  ABS MAX

$-3 \mid -2 \quad 54 \quad 5$   
 $\quad \quad \quad 6 \quad -180$

$-2 \quad 60 \mid -175 = f(-3)$

ABS MIN

$x=0 \quad 0 \mid -2 \quad 54 \quad 5$   
 $\quad \quad \quad 0 \quad 0$

$-2 \quad 54 \mid 5 = f(0)$

$x=4: \quad 4 \mid -2 \quad 54 \quad 5$   
 $\quad \quad \quad -8 \quad 184$

$-2 \quad 46 \mid 184 = f(4)$

$\begin{array}{r} 2 \quad 46 \\ \underline{\quad 4} \\ 184 \end{array}$

S3.1 #s 5, 6, 7, 9, 15-31, 35-41, 47, 53  
DUE THURSDAY

Find Max of  $f(x) = \frac{1}{x}$  on  $[-1, 1]$

There isn't one!

But what about EVT?


Doesn't satisfy continuity requirement on  $[-1, 1]$ .

#42 Find c.v.s

$$g(x) = \sqrt{1-x^2} = (1-x^2)^{\frac{1}{2}}$$

$$y' = \frac{1}{2}(1-x^2)^{-\frac{1}{2}}(-2x)$$

$$= \frac{-x}{\sqrt{1-x^2}}$$



$$x^2 + y^2 = 1$$

$$y^2 = 1 - x^2$$

$$y = \pm \sqrt{1-x^2}$$

$$y' = 0: -x = 0 \Rightarrow x = 0$$

$$y' \neq 0: \sqrt{1-x^2} = 0$$

$$1-x^2 = 0$$

$$x^2 = 1$$

$$x = \pm 1$$

All that was asked

(55)  $f(t) = 2 \cos t + \sin(2t)$  on  $[0, \frac{\pi}{2}]$

$f(0) = 2$   
 $f(\frac{\pi}{2}) = 0$   $\rightarrow$  Min

$f'(t) = -2 \sin t + 2 \cos(2t)$   $\exists \forall x$ , so only looking for  $f'(t) = 0$

$\stackrel{SE}{=} 0 \Rightarrow 2 \cos(2t) - 2 \sin t = 0$


$\Rightarrow \cos(2t) - \sin t = 0$

$\Rightarrow 1 - 2 \sin^2 t - \sin t = 0$

$\Rightarrow 2 \sin^2 t + \sin t - 1 = 0$

$2u^2 + u - 1 = 0$

$(2u - 1)(u + 1) = 0$




$\Rightarrow 2 \sin t = 1$

$\sin t = \frac{1}{2}$

$t = \frac{\pi}{6}$

$\sin t = -1$

Never, on  $[0, \frac{\pi}{2}]$



$f(\frac{\pi}{6}) = 2 \cos(\frac{\pi}{6}) + \sin(2 \cdot \frac{\pi}{6})$

$= 2 \cdot \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{2} = f(\frac{\pi}{6})$



$\rightarrow$  MAX