

---

## Section 1.8 Solutions

---

1. From Definition 1,  $\lim_{x \rightarrow 4} f(x) = f(4)$ .


3. (a)  $f$  is discontinuous at  $-4$  since  $f(-4)$  is not defined and at  $-2$ ,  $2$ , and  $4$  since the limit does not exist (the left and right limits are not the same).

(b)  $f$  is continuous from the left at  $-2$  since  $\lim_{x \rightarrow -2^-} f(x) = f(-2)$ .  $f$  is continuous from the right at  $2$  and  $4$  since

$\lim_{x \rightarrow 2^+} f(x) = f(2)$  and  $\lim_{x \rightarrow 4^+} f(x) = f(4)$ . It is continuous from neither side at  $-4$  since  $f(-4)$  is undefined.

4.  $g$  is continuous on  $[-4, -2)$ ,  $(-2, 2)$ ,  $[2, 4)$ ,  $(4, 6)$ , and  $(6, 8)$ .


7. The graph of  $y = f(x)$  must have a removable discontinuity (a hole) at  $x = 3$  and a jump discontinuity at  $x = 5$ .



$$13. \lim_{x \rightarrow -1} f(x) = \lim_{x \rightarrow -1} (x + 2x^3)^4 = \left( \lim_{x \rightarrow -1} x + 2 \lim_{x \rightarrow -1} x^3 \right)^4 = [-1 + 2(-1)^3]^4 = (-3)^4 = 81 = f(-1).$$

By the definition of continuity,  $f$  is continuous at  $a = -1$ .

17.  $f(x) = \frac{1}{x+2}$  is discontinuous at  $a = -2$  because  $f(-2)$  is undefined.




$$19. f(x) = \begin{cases} 1 - x^2 & \text{if } x < 1 \\ 1/x & \text{if } x \geq 1 \end{cases}$$

The left-hand limit of  $f$  at  $a = 1$  is

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (1 - x^2) = 0. \text{ The right-hand limit of } f \text{ at } a = 1 \text{ is}$$


$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (1/x) = 1. \text{ Since these limits are not equal, } \lim_{x \rightarrow 1} f(x)$$

does not exist and  $f$  is discontinuous at 1.



$$21. f(x) = \begin{cases} \cos x & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ 1 - x^2 & \text{if } x > 0 \end{cases}$$

$\lim_{x \rightarrow 0} f(x) = 1$ , but  $f(0) = 0 \neq 1$ , so  $f$  is discontinuous at 0.



41.


$$f(x) = \begin{cases} 1 + x^2 & \text{if } x \leq 0 \\ 2 - x & \text{if } 0 < x \leq 2 \\ (x - 2)^2 & \text{if } x > 2 \end{cases}$$

$f$  is continuous on  $(-\infty, 0)$ ,  $(0, 2)$ , and  $(2, \infty)$  since it is a polynomial on

each of these intervals. Now  $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (1 + x^2) = 1$  and  $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (2 - x) = 2$ , so  $f$  is

discontinuous at 0. Since  $f(0) = 1$ ,  $f$  is continuous from the left at 0. Also,  $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (2 - x) = 0$ ,

$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x - 2)^2 = 0$ , and  $f(2) = 0$ , so  $f$  is continuous at 2. The only number at which  $f$  is discontinuous is 0.



64.

$$g(x) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ x & \text{if } x \text{ is irrational} \end{cases} \text{ is continuous at 0. To see why, note that } -|x| \leq g(x) \leq |x|, \text{ so by the Squeeze Theorem}$$

$\lim_{x \rightarrow 0} g(x) = 0 = g(0)$ . But  $g$  is continuous nowhere else. For if  $a \neq 0$  and  $\delta > 0$ , the interval  $(a - \delta, a + \delta)$  contains both

infinitely many rational and infinitely many irrational numbers. Since  $g(a) = 0$  or  $a$ , there are infinitely many numbers  $x$  with

$0 < |x - a| < \delta$  and  $|g(x) - g(a)| > |a|/2$ . Thus,  $\lim_{x \rightarrow a} g(x) \neq g(a)$ .