


† Graph of the length of daylight from March 21 through December 21 at various latitudes.




$$L(t) = 2 \sin\left(\frac{2\pi}{365}(t - 80)\right) + 12$$

$$2 \sin\left(\frac{2}{365}\pi(t - 80)\right) + 12$$

31, 20, 21
 $31 + 20 + 21 = 80$, so
 3/21 is $t = 80$.


1.4 #3
 Avg speed is slope



Hours in varied latitudes

	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
30°	12	13.2	13.7	14	13.8	12.8	12	11.2	10.2	10

$T = \text{period} = 365 \text{ days}$
 start @ midline of $y = 12 \text{ hrs of daylight}$.
 Use sine that starts @ $t = 80^{\text{th}}$ day.



$$365 + 80 = 445$$

$$2 \sin(b(x-c)) + d$$

$$= 2 \sin(b(x-80)) + 12$$

Period is 365, so

$$bx = 2\pi \text{ when } x = 365 \Rightarrow$$

$$365b = 2\pi$$

$$b = \frac{2\pi}{365}$$

$$= 2 \sin\left(\frac{2\pi}{365}(x-80)\right) + 12 = \text{HOURS / DAY of sunlight, using March 21st as 1st day.}$$

1.5 - Limit of a Function

1 Intuitive Definition of a Limit Suppose $f(x)$ is defined when x is near the number a . (This means that f is defined on some open interval that contains a , except possibly at a itself.) Then we write

$$\lim_{x \rightarrow a} f(x) = L$$

and say "the limit of $f(x)$, as x approaches a , equals L "

if we can make the values of $f(x)$ arbitrarily close to L (as close to L as we like) by restricting x to be sufficiently close to a (on either side of a) but not equal to a .

Challenge:
make $|f(x) - L|$ small
by making $|x - a|$
small enough!

The limit of the difference quotient is the limit in which we're most interested, and it always takes the form of a 0/0 situation!

Book Example:

$$\lim_{t \rightarrow 0} \frac{\sqrt{t^2+9} - 3}{t^2}$$

$$(a-b)(a+b) = a^2 - b^2$$


$$\frac{\sqrt{t^2+9} - 3}{t^2} = \frac{(\sqrt{t^2+9} - 3)(\sqrt{t^2+9} + 3)}{t^2(\sqrt{t^2+9} + 3)}$$

$$= \frac{t^2+9-9}{t^2(\sqrt{t^2+9} + 3)} = \frac{\cancel{t^2}}{\cancel{t^2}(\sqrt{t^2+9} + 3)} = \frac{1}{\sqrt{t^2+9} + 3} \quad (t \neq 0)$$

$$\xrightarrow{t \rightarrow 0} \frac{1}{\sqrt{9+3}} = \frac{1}{3+3} = \boxed{\frac{1}{6}}$$

Example 4 $f(x) = \sin\left(\frac{\pi}{x}\right)$

$\lim_{x \rightarrow 0} f(x) =$



Using just 0.1, 0.01, 0.001, ...

$$\frac{\pi}{\frac{1}{10}} = 10\pi$$

$$\frac{\pi}{\frac{1}{1000}} = 1000\pi$$



} Integer multiples of π .
 $\sin(n\pi) = 0$.

So the limit is zero if you are too naive. So beware of numerical methods....

EXAMPLE 5 $f(x) = \frac{\sin(x)}{x}$ IMPORTANT! REMEMBER! (S2.4!)

$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$

(Graphical confirmation w/ CAS (Maple))



2 Definition of One-Sided Limits We write

$$\lim_{x \rightarrow a^-} f(x) = L$$

and say the **left-hand limit** of $f(x)$ as x approaches a [or the **limit of $f(x)$ as x approaches a from the left**] is equal to L if we can make the values of $f(x)$ arbitrarily close to L by taking x to be sufficiently close to a with x less than a .

Same thing for right-handed limits. Just turn the "-" into a "+" and replace "left" with "right."

$$\lim_{x \rightarrow a^+} f(x) = L$$


The "regular" limit exists if and only if the left and right limits both exist and both agree.

3 $\lim_{x \rightarrow a} f(x) = L$ if and only if $\lim_{x \rightarrow a^-} f(x) = L$ and $\lim_{x \rightarrow a^+} f(x) = L$

$$\lim_{x \rightarrow 3^-} (x^2 - 5) = 3^2 - 5 = 4$$

$$\lim_{x \rightarrow 3^+} (x^2 - 5) = \text{same.}$$

$$f(x) = \begin{cases} x^2 - 5 & \text{if } x < 0 \\ 2x - 5 & \text{if } x \geq 0 \end{cases}$$



$$\lim_{x \rightarrow 0^-} f(x) = 0^2 - 5 = -5$$

$$\lim_{x \rightarrow 0^+} f(x) = 2(0) - 5 = -5$$


$$\therefore \lim_{x \rightarrow 0} f(x) = -5$$

■ Infinite Limits

EXAMPLE 8 Find $\lim_{x \rightarrow 0} \frac{1}{x^2}$ if it exists.

To indicate the kind of behavior exhibited in Example 8, we use the notation

$$\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$$



This does not mean that we are regarding ∞ as a number. Nor does it mean that the limit exists. It simply expresses the particular way in which the limit does not exist: $1/x^2$ can be made as large as we like by taking x close enough to 0.

In general, we write symbolically

$$\lim_{x \rightarrow a} f(x) = \infty$$

*GIVE ME AN $M > 0$
I can make $f(x) > M$ by
choosing x close enough to a .*

to indicate that the values of $f(x)$ tend to become larger and larger (or “increase without bound”) as x becomes closer and closer to a .

4 Intuitive Definition of an Infinite Limit Let f be a function defined on both sides of a , except possibly at a itself. Then

$$\lim_{x \rightarrow a} f(x) = \infty$$

means that the values of $f(x)$ can be made arbitrarily large (as large as we please) by taking x sufficiently close to a , but not equal to a .

5 Definition Let f be a function defined on both sides of a , except possibly at a itself. Then

$$\lim_{x \rightarrow a} f(x) = -\infty$$

means that the values of $f(x)$ can be made arbitrarily large negative by taking x sufficiently close to a , but not equal to a .


The symbol $\lim_{x \rightarrow a} f(x) = -\infty$ can be read as “the limit of $f(x)$, as x approaches a , is negative infinity” or “ $f(x)$ decreases without bound as x approaches a .” As an example we have

$$\lim_{x \rightarrow 0} \left(-\frac{1}{x^2} \right) = -\infty$$

Useful for a graph, but strictly speaking, these infinite limits do not exist as real numbers.

These infinite limits correspond to vertical asymptotes in graphs.

Sketch the graph of a simple linear/linear rational function.



Section 1.6 - Limit Laws

Limit Laws Suppose that c is a constant and the limits

$$\lim_{x \rightarrow a} f(x) \quad \text{and} \quad \lim_{x \rightarrow a} g(x)$$

exist. Then

$$1. \lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

$$2. \lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$$

$$3. \lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x)$$

$$4. \lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$$

$$5. \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} \quad \text{if } \lim_{x \rightarrow a} g(x) \neq 0$$

§ $c \in \mathbb{R}$, $\lim_{x \rightarrow a} f(x) = L$, $\lim_{x \rightarrow a} g(x) = M$. Then

$$\lim_{x \rightarrow a} (f \pm g) = L \pm M$$

$$\lim_{x \rightarrow a} cf = cL$$

$$\lim_{x \rightarrow a} (fg) = LM$$

$$\lim_{x \rightarrow a} \left(\frac{f}{g}\right) = \frac{L}{M}, \text{ if } M \neq 0.$$

$$\lim_{x \rightarrow a} (f^n) = L^n \quad \forall n \in \mathbb{N}$$

$$\lim_{x \rightarrow a} (c) = c$$

$$\lim_{x \rightarrow a} (x) = a$$

$$\lim_{x \rightarrow a} (x^n) = a^n \quad \forall n \in \mathbb{N}$$

$$\lim_{x \rightarrow a} (\sqrt[n]{f}) = \sqrt[n]{L} \quad \forall n \in \mathbb{N}$$

If $n = 2m$ for $m \in \mathbb{N}$, then assume $L > 0$.

Let's speed the evaluations up:

Direct Substitution Property If f is a polynomial or a rational function and a is in the domain of f , then

$$\lim_{x \rightarrow a} f(x) = f(a)$$

If $f(x) = g(x)$ when $x \neq a$, then $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x)$, provided the limits exist.

This is why there's a string of equal signs from the original difference quotient to the final passage to the limit at the end, after you cancel out the h 's.

