


5-18 Evaluate the surface integral.

7.  $\iint_S yz \, dS,$

$S$  is the part of the plane  $x + y + z = 1$  that lies in the first octant




$$\vec{r} = \langle x, y, 1-x-y \rangle$$

$$\vec{r}_x = \langle 1, 0, -1 \rangle, \quad 1, 0$$

$$\vec{r}_y = \langle 0, 1, -1 \rangle, \quad 0, 1$$

$$\langle 1, 1, 1 \rangle$$

$$\Rightarrow \|\vec{r}_x \times \vec{r}_y\| = \sqrt{3}$$



TI

$$\iint_S yz \, dS = \int_0^1 \int_0^{1-x} y(1-x-y) \sqrt{3} \, dy \, dx$$

$$\sqrt{3} \int_0^1 \int_0^{1-x} [y - xy - y^2] \, dy \, dx = \sqrt{3} \int_0^1 \left[ \frac{1}{2}y^2 - \frac{1}{2}y^2x - \frac{1}{3}y^3 \right]_0^{1-x} dx$$

$$= \frac{\sqrt{3}}{2} \int_0^1 (1-x)^2 - x(1-x)^2 \, dx - \frac{\sqrt{3}}{3} \int_0^1 (1-x)^3 \, dx$$

$$= \frac{\sqrt{3}}{2} \int_0^1 (x^2 - 2x + 1 - x^3 + 2x^2 - x) \, dx + \frac{\sqrt{3}}{3} \left[ \frac{(1-x)^4}{4} \right]_0^1$$

$$= \frac{\sqrt{3}}{2} \int_0^1 (-x^3 + 3x^2 - 3x + 1) \, dx + \frac{\sqrt{3}}{3} \left[ -\frac{1}{4} \right]$$

$$= \frac{\sqrt{3}}{2} \left[ -\frac{x^4}{4} + x^3 - \frac{3}{2}x^2 + x \right]_0^1 - \frac{\sqrt{3}}{12}$$

$$= \frac{\sqrt{3}}{2} \left[ -\frac{1}{4} + 1 - \frac{3}{2} + 1 \right] - \frac{\sqrt{3}}{12}$$

$$= \frac{\sqrt{3}}{2} \left[ 2 - \frac{7}{4} \right] - \frac{\sqrt{3}}{12} = \frac{-\sqrt{3}}{2} \left[ \frac{8-7}{4} \right] - \frac{\sqrt{3}}{12}$$

$$= \frac{+\sqrt{3}}{8} - \frac{\sqrt{3}}{12} = \frac{+3\sqrt{3}}{24} - \frac{2\sqrt{3}}{24} = \boxed{\frac{\sqrt{3}}{24}}$$

9.  $\iint_S yz \, dS,$


$S$  is the surface with parametric equations  $x = u^2, y = u \sin v,$   
 $z = u \cos v, 0 \leq u \leq 1, 0 \leq v \leq \pi/2$

14.  $\iint_S y^2 dS,$

$S$  is the part of the sphere  $x^2 + y^2 + z^2 = 4$  that lies inside the cylinder  $x^2 + y^2 = 1$  and above the  $xy$ -plane

17.  $\iint_S (z + x^2 y) dS,$

$S$  is the part of the cylinder  $y^2 + z^2 = 1$  that lies between the planes  $x = 0$  and  $x = 3$  in the first octant



$$x = x, y = \cos \theta, z = \sin \theta$$

$$0 \leq \theta \leq \frac{\pi}{2}, 0 \leq x \leq 3$$

$$\vec{r} = \langle x, \cos \theta, \sin \theta \rangle$$

$$\vec{r}_x = \langle 1, 0, 0 \rangle, \quad 1, \quad 0$$

$$\vec{r}_\theta = \langle 0, -\sin \theta, \cos \theta \rangle, \quad 0, \quad -\sin \theta$$

$$\langle 0, -\cos \theta, -\sin \theta \rangle$$

$$\Rightarrow \|\vec{r}_x \times \vec{r}_\theta\| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1$$

$$\iint_S (z + x^2 y) dS$$

$$\iint_S (z + x^2 y) dS = \int_0^3 \int_0^{\frac{\pi}{2}} (\sin \theta + x^2 \cos \theta) d\theta dx$$

$$= \int_0^3 \left[ -\cos \theta + x^2 \sin \theta \right]_0^{\frac{\pi}{2}} dx$$


$$= \int_0^3 (1 + x^2) dx = \left[ x + \frac{x^3}{3} \right]_0^3 = 3 + 9 = 12$$

41. A fluid has density  $870 \text{ kg/m}^3$  and flows with velocity  $\mathbf{v} = z \mathbf{i} + y^2 \mathbf{j} + x^2 \mathbf{k}$ , where  $x$ ,  $y$ , and  $z$  are measured in meters and the components of  $\mathbf{v}$  in meters per second. Find the rate of flow outward through the cylinder  $x^2 + y^2 = 4$ ,  $0 \leq z \leq 1$ .

$$\int_S \mathbf{F} \cdot d\mathbf{S} = \int_S \mathbf{F} \cdot \mathbf{n} \, dS$$

$$= \int_S \mathbf{F} \cdot (\mathbf{r}_u \times \mathbf{r}_v) \, dA$$

$$\mathbf{v} = \langle z, y^2, x^2 \rangle, \quad x, y, z \sim \text{m}, \quad \mathbf{v} \sim \text{m/s}, \quad S = \{(x, y, z) \mid x^2 + y^2 = 4, 0 \leq z \leq 1\}$$



$$S_1 = \text{Bottom}, \quad S_2 = \text{Top}, \quad S_3 = \text{sides}$$

$$\text{Bottom: } \mathbf{r}_1 = \langle 0, 0, -1 \rangle \quad (\text{But lets work it out like a machine.})$$

$$0 \leq r \leq 2, \quad 0 \leq \theta \leq 2\pi, \quad z = 0!$$

$$\mathbf{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, 0 \rangle$$

$$\mathbf{r}_r = \langle \cos \theta, \sin \theta, 0 \rangle, \quad \mathbf{r}_\theta = \langle -r \sin \theta, r \cos \theta, 0 \rangle$$

$$\mathbf{r}_r \times \mathbf{r}_\theta = \langle -r \sin \theta, r \cos \theta, 0 \rangle \times \langle \cos \theta, \sin \theta, 0 \rangle$$

$$= \langle 0, 0, r \cos^2 \theta + r \sin^2 \theta \rangle$$

$$= \langle 0, 0, r \rangle \quad \text{But oriented down.}$$

$$\mathbf{v} = \langle z, y^2, x^2 \rangle$$

$$= \langle 0, r^2 \sin^2 \theta, r^2 \cos^2 \theta \rangle$$

$$\mathbf{v} \cdot (\mathbf{r}_r \times \mathbf{r}_\theta) =$$

$$\langle 0, r^2 \sin^2 \theta, r^2 \cos^2 \theta \rangle \cdot \langle 0, 0, r \rangle = -r^3 \cos^2 \theta$$

$$\int_{S_1} \text{so, } -\rho \int_0^{2\pi} \int_0^2 r^3 \cos^2 \theta \, dr \, d\theta = -\rho \int_0^{2\pi} \left( \frac{1 + \cos 2\theta}{2} \right) d\theta \int_0^2 r^3 \, dr \quad \text{etc.}$$

$$S_2: \mathbf{r}_2(r, \theta) = \langle r \cos \theta, r \sin \theta, 1 \rangle$$

$$\mathbf{r}_r = \langle \cos \theta, \sin \theta, 0 \rangle, \quad \mathbf{r}_\theta = \langle -r \sin \theta, r \cos \theta, 0 \rangle$$

$$\mathbf{r}_r \times \mathbf{r}_\theta = \langle -r \sin \theta, r \cos \theta, 0 \rangle \times \langle \cos \theta, \sin \theta, 0 \rangle$$

$$\langle 0, 0, r \cos^2 \theta + r \sin^2 \theta \rangle = \langle 0, 0, r \rangle \quad \frac{4\rho}{2}$$

$$\vec{v} \cdot (\vec{r}_r \times \vec{r}_\theta) = \langle 0, r^2 \sin^2 \theta, r^2 \cos^2 \theta \rangle \cdot \langle 0, 0, r \rangle$$

$$= r^3 \cos^2 \theta$$

$$\int_2^1: \int_0^{2\pi} \int_0^1 r^3 \cos^2 \theta \, dr \, d\theta$$

$\int_1 + \int_2 = 0!$  (opposite signs from opposite orientation)

$$\int_3^1 \text{ sides: } \vec{r}_3(\theta, z) = \langle 2 \cos \theta, 2 \sin \theta, z \rangle$$

$$\vec{r}_{3\theta} = \langle -2 \sin \theta, 2 \cos \theta, 0 \rangle \quad -2 \sin \theta, 2 \cos \theta$$

$$\vec{r}_{3z} = \langle 0, 0, 1 \rangle \quad 0, 0$$

$$\langle 2 \cos \theta, 2 \sin \theta, 0 \rangle = \vec{r}_{3\theta} \times \vec{r}_{3z}$$

$$\vec{v}(\theta, z)$$

$$= \langle z, 4 \sin^2 \theta, 4 \cos^2 \theta \rangle \quad \vec{v} = \langle z, y^2, x^2 \rangle$$

$$\vec{v} \cdot (\vec{r}_{3\theta} \times \vec{r}_{3z}) = \langle z, 4 \sin^2 \theta, 4 \cos^2 \theta \rangle \cdot \langle 2 \cos \theta, 2 \sin \theta, 0 \rangle$$

$$= 2z \cos \theta + 8 \sin^2 \theta \cos \theta + 0$$

$$\text{So, } \int_0^1 \int_0^{2\pi} (2z \cos \theta + 8 \sin^2 \theta \cos \theta) \, d\theta \, dz = 0$$

So Flux is zero!

$$= \int_0^1 \left[ 2z \sin \theta + 8 \frac{\sin^3 \theta}{3} \right]_0^{2\pi} dz = \int_0^1 0 \, dz = 0$$