

16.9 Change of Variable(s) in Multiple Integrals

Recall the Substitution Rule: THIS FIRST PAGE OF NOTES IS MESSED-UP. DON'T KNOW WHAT I WAS THINKING. WILL BE RE-WORKED, MONDAY...

$$\int_a^b f(x) dx = \int_c^d f(g(u))g'(u) du$$

$$\int_c^d f(u) du = \int_a^b f(g(x))g'(x) dx$$

$x = g(u)$ and $a = g(c), b = g(d)$


$$\int_a^b f(x) dx = \int_c^d f(x(u)) \frac{dx}{du} du$$

$$\int_0^2 \cos(u) du = \int_0^1 \cos(2x) \cdot 2 dx$$

$u = g(x) = 2x$
 $du = g'(x) dx = 2 dx$

Example: Trigonometric Substitution to simplify the integration process.

Find the area inside half a circle of radius $r = 1$.



$$x^2 + y^2 = 1$$

$$y^2 = 1 - x^2$$

$$y = \pm \sqrt{1 - x^2}$$

$y = \sqrt{1 - x^2}$ is the top


$$\text{Area} = \int_{-1}^1 \sqrt{1-x^2} dx$$

$x = 0$
 $u = 2x = 0 = g(0)$
 $x = 1$
 $u = 2 \cdot 1 = 2 = g(1)$

$$\sqrt{1-x^2} = \sqrt{1 - (\cos \theta)^2}$$

$$= \sqrt{1 - \cos^2 \theta} = \sqrt{\sin^2 \theta}$$

$$= |\sin \theta| = \sin \theta \quad \forall \theta \in [0, \pi]$$



$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$x = \cos \theta$$

$$dx = -\sin \theta d\theta$$

$$x = -1 = \cos \theta$$

$$\theta = \pi$$

$$x = 1 = \cos \theta$$

$$\theta = 0$$

$$\int_{\pi}^0 \sin \theta (-\sin \theta d\theta)$$

$$= - \int_{\pi}^0 \sin^2 \theta d\theta$$

$$= \int_0^{\pi} \frac{1 - \cos(2\theta)}{2} d\theta$$


$$= \int_0^{\pi} \frac{1}{2} d\theta - \frac{1}{2} \cdot \frac{1}{2} \int_0^{\pi} \cos(2\theta) \cdot 2 d\theta$$

$$= \frac{1}{2}\pi - \frac{1}{4} [\sin 2\theta]_0^{\pi} = \frac{\pi}{2}$$

We've already extended this to double integrals and polar coordinates:

$$x = r \cos \theta$$

$$y = r \sin \theta$$



$$\iint_R f(x, y) dA = \iint_S f(r \cos \theta, r \sin \theta) r dr d\theta$$

is how we go from
 $dA = dx dy = dy dx$
to
 $r dr d\theta$

$$x = r \cos \theta, y = r \sin \theta$$

JACOBIAN IS THE BRIDGE

$$T(r, \theta) = (x, y)$$


The idea will be to

Do a T^{-1} :

$$T(u, v) = (x, y)$$

$$T^{-1}(T(r, \theta)) = T^{-1}(x, y) = (r, \theta)$$

$$x = g(u, v)$$




$$y = h(u, v)$$

We usually assume that T is a C^1 transformation, which means that g and h have continuous first-order partial derivatives.

Transformations (Mappings) from one domain *onto* another.

If $T(u_1, v_1) = (x_1, y_1)$, then the point (x_1, y_1) is called the **image** of the point (u_1, v_1)



$$u = G(x, y)$$

$$v = H(x, y)$$



EXAMPLE 1 A transformation is defined by the equations

$$T^{-1}(x, y) = (u^2 - v^2, 2uv) \quad x = u^2 - v^2 \quad y = 2uv$$

Find the image of the square $S = \{(u, v) \mid 0 \leq u \leq 1, 0 \leq v \leq 1\}$

SOLUTION The transformation maps the boundary of S into the boundary of the image.

is reported as a fact, with no real proof, even though it is true.



$$T(S_1) : v = 0, 0 \leq u \leq 1$$

$$x = u^2 - v^2 = u^2 \quad y = 2uv = 0$$

$$\Rightarrow 0 \leq x \leq 1$$

$$T(S_2) : u = 1, 0 \leq v \leq 1$$

$$x = 1 - v^2, \quad y = 2v \Rightarrow v = \frac{1}{2}y$$


$$x = 1 - v^2 = 1 - \left(\frac{1}{2}y\right)^2 = 1 - \frac{1}{4}y^2$$

$$0 \leq v \leq 1$$

$$\& y = 2v \Rightarrow$$

$$0 \leq 2v \leq 2$$

$$0 \leq y \leq 2$$



$\iint_R f(x,y) dA$
 \downarrow
 $\iint_S f(u,v) dA$ *in terms of u, v.*

$\mathbf{r}(u_0, v_0)$
 $\mathbf{r}(u_0, v_0 + \Delta v)$
 $\mathbf{r}(u_0 + \Delta u, v_0)$
 $\mathbf{r}(u_0 + \Delta u, v_0 + \Delta v)$

$\mathbf{r}(u_0, v_0)$
 $\mathbf{r}(u_0 + \Delta u, v_0)$
 $\mathbf{r}(u_0, v_0 + \Delta v)$
 $\mathbf{r}(u_0 + \Delta u, v_0 + \Delta v)$

$\Delta u \mathbf{r}_u$
 $\Delta v \mathbf{r}_v$

Tangent Line Segment to the boundary @ $(u_0, v_0) = \mathbf{r}(u_0, v_0)$

Derivation:
 $\Delta \bar{r} = \bar{b} = \mathbf{r}(u_0, v_0 + \Delta v) - \mathbf{r}(u_0, v_0)$
 $\Rightarrow \frac{\Delta \bar{r}}{\Delta v} = \frac{\mathbf{r}(u_0, v_0 + \Delta v) - \mathbf{r}(u_0, v_0)}{\Delta v} \approx \frac{d\bar{r}}{dv} = \mathbf{r}_v$
 $\Rightarrow \bar{b} = \mathbf{r}(u_0, v_0 + \Delta v) - \mathbf{r}(u_0, v_0)$
 $= \frac{\mathbf{r}(u_0, v_0 + \Delta v) - \mathbf{r}(u_0, v_0)}{\Delta v} \cdot \Delta v$
 $\approx \frac{d\bar{r}}{dv} \cdot \Delta v = \mathbf{r}_v \Delta v$
 $\bar{a} = \frac{d\bar{r}}{du} \cdot \Delta u = \mathbf{r}_u \Delta u$
 $\Rightarrow \|\bar{a} \times \bar{b}\| \approx \|(\mathbf{r}_u \Delta u) \times (\mathbf{r}_v \Delta v)\| = \|\mathbf{r}_u \times \mathbf{r}_v\| \Delta u \Delta v = dA$
The increment of area for the conversion!

$$\iint_R f(x, y) dA = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

7 DEFINITION The **Jacobian** of the transformation T given by $x = g(u, v)$ and $y = h(u, v)$ is


$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

↳ Area of the region R using mapping T & domain S !

Double Integral in Polar Coordinates. Where did the $r \, dr \, d\theta$ come from?

We know, but here we do it in terms of the new machinery:

$$\int_0^1 \int_0^{\sqrt{4-x^2}} (x^2 + y^2) \, dy \, dx$$



$$x = r \cos \theta, \quad y = r \sin \theta$$

$$T(r \cos \theta, r \sin \theta) = (x, y)$$

$$\vec{r}(r, \theta) = \langle r \cos \theta, r \sin \theta \rangle$$

$$\vec{r}_r = \langle \cos \theta, \sin \theta \rangle$$

$$\vec{r}_\theta = \langle -r \sin \theta, r \cos \theta \rangle$$

$$\|\vec{r}_r \times \vec{r}_\theta\|$$


$$\begin{vmatrix} \cos \theta & \sin \theta \\ -r \sin \theta & r \cos \theta \end{vmatrix} = |r \cos^2 \theta + r \sin^2 \theta|$$

$$= |r| = r \quad \text{if } 0 \leq r$$

It's the r in the $r \, dr \, d\theta$!

#12

$\iint_R (4x+8y) dA$, where R is the parallelogram with corners $(-1,3), (1,-3), (3,-1), (1,5)$




$$\vec{r}(u,v) = \left\langle \frac{1}{4}u + \frac{1}{4}v, \frac{1}{4}v - \frac{3}{4}u \right\rangle$$

$$\vec{r}_u = \left\langle \frac{1}{4}, -\frac{3}{4} \right\rangle, \vec{r}_v = \left\langle \frac{1}{4}, \frac{1}{4} \right\rangle$$

$$\begin{vmatrix} \frac{1}{4} & -\frac{3}{4} \\ \frac{1}{4} & \frac{1}{4} \end{vmatrix} = \frac{1}{16} + \frac{3}{16} = \frac{4}{16} = \frac{1}{4} = \frac{2(x,y)}{2(u,v)}$$

$$4x + 8y = 4\left(\frac{1}{4}u + \frac{1}{4}v\right) + 8\left(\frac{1}{4}v - \frac{3}{4}u\right) = u + v + 2v - 6u = -5u + 3v$$



To get T^{-1} , solve for u and v :

$$\begin{aligned} x &= \frac{1}{4}u + \frac{1}{4}v \\ y &= \frac{1}{4}v - \frac{3}{4}u \end{aligned}$$

$$\begin{aligned} \frac{1}{4}u + \frac{1}{4}v &= x \\ -\frac{3}{4}u + \frac{1}{4}v &= y \end{aligned} \Rightarrow \left[\begin{array}{cc|c} \frac{1}{4} & \frac{1}{4} & x \\ -\frac{3}{4} & \frac{1}{4} & y \end{array} \right]$$

T

Reduced Row-Echelon Form: $\left[\begin{array}{cc|c} 1 & 0 & x-y \\ 0 & 1 & 3x+y \end{array} \right]$

T^{-1} \Rightarrow $\begin{cases} u = x - y \\ v = 3x + y \end{cases}$

$(-1,3), (1,-3), (3,-1), (1,5)$

$$T^{-1}(x,y) = (u,v) = (x-y, 3x+y)$$

$$T^{-1}(-1,3) = (-1-3, 3(-1)+3) = (-4,0)$$

$$T^{-1}(1,3) = (1-3, 3(1)-3) = (-2,0)$$

$$T^{-1}(3,-1) = (3-1, 3(3)+(-1)) = (2,10) \quad \text{No. } (3 - (-1), 3(3) + (-1)) = (4, 8)$$

$$T^{-1}(1,5) = (1-5, 3(1)+5) = (-4,8)$$

